



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

*Reports of States and yearly and monthly reports of cities.*

CONNECTICUT.—Month of November, 1896. Reports to the State board of health from 165 towns, having an aggregate population of 852,540, show a total of 983 deaths, including phthisis pulmonalis, 105; enteric fever, 24; scarlet fever, 6; measles, 4; diphtheria and croup, 54, and whooping cough, 3.

MARYLAND—*Baltimore*.—Month of November, 1896. Estimated population, white, 431,054; colored, 75,344; total, 506,398. Deaths, white, 482; colored, 152; total, 634, including phthisis pulmonalis, 83; enteric fever, 22; scarlet fever, 1; diphtheria, 31, and croup, 2.

MICHIGAN.—Week ended December 19, 1896. Reports to the State board of health from 85 observers indicate that whooping cough, inflammation of brain, scarlet fever, diarrhea, remittent fever increased, and inflammation of bowels and consumption decreased in area of prevalence.

Phthisis pulmonalis was reported present during the week at 183 places, measles at 17, scarlet fever at 50, diphtheria at 56, enteric fever at 32, and whooping cough at 19.

MISSOURI.—Month of November, 1896. Estimated population, 165,000. Total deaths, 142, including phthisis pulmonalis, 10; enteric fever, 2; diphtheria, 9; croup, 4; measles 1, and whooping cough, 1.

NEBRASKA—*Omaha*.—Month of October, 1896. Estimated population, 140,452. Total deaths, 95, including phthisis pulmonalis, 10; enteric fever, 5; scarlet fever, 1, and diphtheria and croup, 10.

Month of November, 1896. Total deaths, 94, including phthisis pulmonalis, 8; enteric fever, 1; scarlet fever, 1, and diphtheria and croup, 7.

NEW YORK—*Yonkers*.—Month of November, 1896. Estimated population, 38,000. Total deaths, 44, including phthisis pulmonalis, 4, and diphtheria, 3.

OHIO—*Columbus*.—Month of November, 1896. Estimated population, 130,000. Total deaths, 121, including phthisis pulmonalis, 15; enteric fever, 9, and diphtheria, 12.

PENNSYLVANIA—*Westchester*.—Month of November, 1896. Estimated population, 10,000. Total deaths, 16, including phthisis pulmonalis, 1, and diphtheria, 1.

UTAH—*Salt Lake City*.—Month of November, 1896. Estimated population, 70,000. Total deaths, 47, including phthisis pulmonalis, 5; enteric fever, 5; scarlet fever, 1, and diphtheria, 1.